



SHORT GENOME REPORT

Open Access

Complete genome sequence of *Salmonella enterica* subspecies *arizonae* str. RKS2983

Chun-Xiao Wang¹, Song-Ling Zhu¹, Xiao-Yu Wang¹, Ye Feng², Bailiang Li¹, Yong-Guo Li³, Randal N Johnston⁴, Gui-Rong Liu¹, Jin Zhou^{5*} and Shu-Lin Liu^{1,3,6*}

Abstract

Salmonella arizonae (also called *Salmonella* subgroup IIIa) is a Gram-negative, non-spore-forming, motile, rod-shaped, facultatively anaerobic bacterium. *S. arizonae* strain RKS2983 was isolated from a human in California, USA. *S. arizonae* lies somewhere between *Salmonella* subgroups I (human pathogens) and V (also called *S. bongori*; usually non-pathogenic to humans) and so is an ideal model organism for studies of bacterial evolution from non-human pathogen to human pathogens. We hence sequenced the genome of RKS2983 for clues of genomic events that might have led to the divergence and speciation of *Salmonella* into distinct lineages with diverse host ranges and pathogenic features. The 4,574,836 bp complete genome contains 4,203 protein-coding genes, 82 tRNA genes and 7 rRNA operons. This genome contains several characteristics not reported to date in *Salmonella* subgroup I or V and may provide information about the genetic divergence of *Salmonella* pathogens.

Keywords: *S. enterica* subspecies *arizonae* RKS2983, Facultative anaerobe, Genomic evolution, Host-adapted, *Salmonella* pathogenicity islands

Introduction

Salmonella are Gram-negative facultative anaerobic bacteria of the family *Enterobacteriaceae* inhabiting the gastrointestinal tract of a wide variety of animals. There are currently over 2,600 serotypes (also called serovars) documented in the genus *Salmonella*. By chromosomal DNA hybridization experiments and MLEE, *Salmonella* currently are classified into two species, *S. enterica* and *S. bongori* (formerly subgroup V). The species *S. enterica* is further divided into six subspecies, including *S. enterica* subspecies *enterica*, *S. enterica* subspecies *salamae*, *S. enterica* subspecies *arizonae*, *S. enterica* subspecies *diarizonae*, *S. enterica* subspecies *houtenae*, and *S. enterica* subspecies *indica*, corresponding to the former subgroups I, II, IIIa, IIIb, IV and VI, respectively. Additionally, subgroup VII was described by Boyd et al. [1,2]. *Salmonella*

taxonomy is a dynamic field of research and many issues remain unsolved, especially regarding species definition [3-5]. To avoid confusions, therefore, we use the traditional *Salmonella* classification system and the terms subgroup and serotype rather than subspecies or serovar (see more detailed explanation in [5]). Most of *Salmonella* infections in warm-blooded animals are caused by *Salmonella* subgroup I serotypes, and non-subgroup I serotypes are typically associated with cold-blooded vertebrates and rarely colonize the intestines of warm-blooded animals.

Salmonella evolved from a common ancestor with *Escherichia coli* about 120–150 million years ago [6,7]. During the evolutionary process, several key genomic events might have led bacteria to diverge, such as gene mutation and gene acquisition or loss [8]. Importantly, numerous lines of evidence have indicated that gene acquisition and loss are the major force driving the

* Correspondence: jinzhouh85@163.com; slliu@ucalgary.ca⁵Department of Hematology, The First Affiliated Hospital, Harbin Medical University, Harbin, China¹Genomics Research Center, Harbin Medical University, Harbin, China

Full list of author information is available at the end of the article

evolution of virulence in *Salmonella* [9]. In fact, it has been postulated that the evolution of *Salmonella*-specific virulence can be divided into three phases. The first phase is the split of *Salmonella* and *E. coli*

Table 1 Classification and general features of *S. arizonae* RKS2983

MIGS ID	Property	Term	Evidence code ^a
	Current classification	Domain <i>Bacteria</i>	TAS [34]
		Phylum <i>Proteobacteria</i>	TAS [35]
		Class <i>Gammaproteobacteria</i>	TAS [35,36]
		Order " <i>Enterobacteriales</i> "	TAS [37-39]
		Family <i>Enterobacteriaceae</i>	TAS [39,40]
		Genus <i>Salmonella</i>	TAS [40-41]
		Species <i>Salmonella enterica</i>	TAS [41,42]
		Subspecies <i>Salmonella enterica</i> subsp. <i>arizonae</i>	TAS [42]
		Strain RKS2983	TAS [42]
		Serovar 62:z36:-	TAS [42]
	Gram stain	Negative	IDA
	Cell shape	Rod-shaped	IDA
	Motility	Motile	IDA
	Sporulation	Non-sporulating	IDA
	Temperature range	Mesophilic	IDA
	Optimum temperature	35°C–37°C	IDA
	pH	7.2–7.6	IDA
	Carbon source	Glucose	IDA
MIGS-6	Habitat	Human	TAS [42]
MIGS-6.3	Salinity	Medium	IDA
MIGS-22	Oxygen requirement	Facultative anaerobes	IDA
MIGS-15	Biotic relationship	Endophyte	IDA
MIGS-14	Pathogenicity	Pathogenic	IDA
MIGS-4	Geographic location	California, USA	TAS [42]
MIGS-5	Sample collection time	1985	TAS [42]
MIGS-4.1	Latitude	Not report	NAS
MIGS-4.2	Longitude	Not report	NAS
MIGS-4.3	Depth	Not report	NAS
MIGS-4.4	Altitude	Not report	NAS

a.) Evidence codes - IDA: Inferred from Direct Assay; TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [43].

by the *Salmonella* acquisition of *Salmonella* pathogenicity island 1, which is present in all lineages of *Salmonella* but absent from *E. coli*. SPI-1 encodes virulence factors that strengthen the infection of *Salmonella* serotypes by different mechanisms, including the invasiveness of the bacteria into intestinal epithelial cells [10], induction of neutrophil recruitment, and secretion of intestinal fluid [11-13]. The second phase is the divergence of *Salmonella* into *S. bongori* and *S. enterica*; this pathogenic lineage acquired SPI-2 [14-17], which contains genes encoding a type III secretion system that is required for survival in macrophages [18]. The third phase is the adaptation of *Salmonella* subgroup I to warm-blooded animals, but the key genomic events involved remain unknown.

Genome sequencing efforts in *Salmonella* have mostly focused on *Salmonella* subgroup I serotypes, largely due to their pathogenicity in humans. In this study, we sequenced the genome of a strain from *Salmonella* subgroup IIIa (also known as *Salmonella arizonae*), which lies somewhere between *Salmonella* subgroups I and V in evolution. Based on the important evolutionary position of *Salmonella* subgroup IIIa, we anticipated that its genomic comparisons with other *Salmonella* subgroups, especially subgroups I and V, may provide novel insights into the evolutionary transition of *Salmonella* adaptation from cold- to warm-blooded hosts.

Organism information

Classification and features

S. arizonae is classified to Class *Gammaproteobacteria*, Order *Enterobacteriales*, Family *Enterobacteriaceae* and

Table 2 Project information

MIGS ID	Property	Term
MIGS-31	Finishing quality	Finished
MIGS-28	Libraries used	Illumina Paired-End library and SOLiD mate_pair library (2 x 50 bp)
MIGS-29	Sequencing platforms	Illumina HiSeq 2000 and SOLiD 3.0
MIGS-31.2	Fold coverage	100 x
MIGS-30	Assemblers	SOAPdenovo v1.05
MIGS-32	Gene calling method	Glimmer software that used in the RAST pipeline
MIGS 13	Genbank ID	CP006693.1
	Genbank date of release	September 22, 2014
	GOLD ID	GI686507741
	BIOPROJECT	PRJNA215272
	Source material identifier	CDC 409-85
	Project relevance	Evolution in bacteria




Figure 1 Graphical circular map of the *S. arizonae* RKS2983 genome. From the outside to the center: genes on forward strand (color by COG categories), genes on reverse strand (color by COG categories), GC content, and GC skew. The map was generated with the CGviewer software.

Genus *Salmonella* (Table 1). *S. arizonae* was first described in 1939 by the name *Salmonella dar es salaam* and was categorized as *Salmonella* subgroup IIIa, later named *S. arizonae* [19]. *S. arizonae* is a rare cause of human infection and is naturally found in reptiles.

We obtained RKS2983 from the *Salmonella* Genetic Stock Center (SGSC) as one of the strains in the set of *Salmonella* Reference Collection C strain (SARC6) [2]; it was initially isolated from a human of California in 1985. It is, like other *Salmonella* bacteria, Gram-negative with diameters around 0.7 to 1.5 µm and lengths of 2 to 5 µm, facultatively anaerobic, non-spore-forming, and predominantly motile with peritrichous flagella. The bacteria were grown at 37°C in Luria broth with pH of 7.2-7.6. Detailed information on the strain can be found at SGSC [20].

Genome sequencing information

Genome project history

This complete genome project was deposited in the Genomes On-Line Database (GOLD) and the complete genome sequence of strain RKS2983 was deposited at DDBJ/EMBL/GenBank under the accession CP006693.1.

Table 2 presents the project information and its association with MIGS version 2.0 [21].

Growth conditions and DNA isolation

S. arizonae RKS2983 was cultured to mid-logarithmic phase in 50 ml of Luria Broth on a gyratory shaker at

Table 3 Nucleotide content and gene count levels of the genome

Attribute	Value	% of total ^a
Genome Size (bp)	4,574,836	
G + C content (bp)	2,356,040	51.50
Coding region (bp)	3,924,843	85.79
Total genes ^b	4,390	
rRNA genes	22	0.50
tRNA genes	82	1.87
Protein-coding genes	4,203	95.70
Pseudogenes	98	2.23
Frameshifted Genes	78	1.78
Genes assigned to COGs	3,383	77.06

a.) The total is based on either the size of the genome in base pairs or the total number of protein coding genes in the annotated genome.

37°C. DNA was isolated from the cells using a CTAB bacterial genomic DNA isolation method [22].

Genome sequencing and assembly

The genome of *S. arizonae* RKS2983 was sequenced by use of two sequencing platforms, SOLiD 3.0 and Illumina HiSeq 2000. First, genomic DNA was sequenced with the Illumina sequencing platform by the paired-end strategy (2×100 bp) and the details of library construction and sequencing can be found at the Illumina web site [23]. The sequence data from Illumina HiSeq 2000 were assembled by SOAPdenovo v1.05 and the assembly contained 103 scaffolds with a genome size of 4.5 Mb. Then, the genomic DNA was sheared into 3 kb fragments by the Hydroshear instrument and was sequenced on a SOLiD sequencer by the mate-pair strategy (2 × 50 bp) according to the manual for the instrument (Applied Biosystems). The two sets of data from different methods were assembled by the velvet v1.2.09 software. The final assembly contained 20 scaffolds. Gaps between contigs were closed by PCR amplification using ABI3730 sequencer.

Genome annotation

Genes were predicted by Rapid Annotation using Subsystem Technology [24] with Glimmer 3 [25] followed by manual curation. The predicted coding sequences (CDSs) were translated and used to search the National Center for Biotechnology Information non-redundant database and Clusters of Orthologous Groups databases. These data sources were combined to assert a product description for each predicted protein. Then, we compared them with the annotated genes from four available *Salmonella* genomes, including *S. typhi* Ty2, *S. typhimurium* LT2 (AE006468) [26], *S. arizonae* RKS2980 (CP000880) [27] and *S. bongori* NCTC12419 (NC_015761) [17]. Non-coding genes and miscellaneous features were predicted using tRNAscanSE [28], RNAMMer [29], Rfam [30] and TMHMM [31].

Genome properties

The genome (Figure 1) consists of a chromosome of 4,574,836 bp (51.5% GC content) with 4,390 genes predicted, including 4,203 protein-coding genes, 22 rRNA genes, 82 tRNA genes and 98 pseudogenes. The properties and the statistics of the genome are summarized in Tables 3 and 4.

Insights from the genome sequence

We first looked into the genetic relatedness of *Salmonella* and *E. coli*. For this, we concatenated the 945 genes common to the 25 sequenced strains analyzed in this

Table 4 Number of genes associated with the 25 general COG functional categories

Code	Value	% of total ^a	Description
J	168	3.83	Translation, ribosomal structure and biogenesis
A	1	0.02	RNA processing and modification
K	0	0.00	Transcription
L	216	4.92	Replication, recombination and repair
B	0	0.00	Chromatin structure and dynamics
D	32	0.73	Cell cycle control, mitosis and meiosis
Y	0	0.00	Nuclear structure
V	44	1.00	Defense mechanisms
T	106	2.41	Signal transduction mechanisms
M	223	5.08	Cell wall/membrane biogenesis
N	89	2.03	Cell motility
Z	0	0.00	Cytoskeleton
W	0	0.00	Extracellular structures
U	44	1.00	Intracellular trafficking and secretion
O	137	3.12	Posttranslational modification, protein turnover, chaperones
C	240	5.47	Energy production and conversion
G	307	6.99	Carbohydrate transport and metabolism
E	314	7.15	Amino acid transport and metabolism
F	76	1.73	Nucleotide transport and metabolism
H	142	3.23	Coenzyme transport and metabolism
I	88	2.00	Lipid transport and metabolism
P	182	4.15	Inorganic ion transport and metabolism
Q	53	1.21	Secondary metabolites biosynthesis, transport and catabolism
R	311	7.08	General function prediction only
S	348	7.93	Function unknown
-	1007	22.94	Not in COGs

a.) The total is based on the total number of protein coding genes in the annotated genome.

study and conducted comparisons using BLAST with the parameters set at >70% DNA identity and >0.7 gene length ratio to categorize genes into common genes. The multiple sequence alignment program MAFFT program [32] was used to align the gene sequences of the *Salmonella* and *E. coli* strains. Phylogenetic trees were constructed with the aligned gene sequences using the Neighbor-Joining methods based on 1,000 randomly selected bootstrap replicates by MEGA 4.0 software [33]. The tree showed that *S. bongori* positioned between *Salmonella* subgroup I and *E. coli*, *S. arizonae* RKS2983 positioned between *Salmonella* subgroup I and *S. bongori*, and all *Salmonella* subgroup I strains were clustered together (Figure 2).




Figure 2 Phylogenetic tree highlighting the position of *S. arizonae* RKS2983 (shown in bold) relative to strains of other *Salmonella* lineages. The corresponding GenBank accession numbers are displayed in parentheses. The tree was built based on the comparison of concatenated nucleotide sequences of 945 conserved genes in all strains. Individual orthologous sequences were aligned by the MAFFT program [32] and concatenated. The phylogenetic tree was constructed by using the MEGA 4.0 software [33] with Neighbor-Joining method. The bootstrap values are shown at branch points.




Figure 3 Venn diagram showing the core genes in *S. arizonae* RKS2983, *S. bongori* NCTC 12419 and *S. typhimurium* LT2. The core genes conducted using BLAST with the parameters set at ">70% DNA identity and >0.7 gene length ratio".

Table 5 Distribution of known SPIs in four representation genomes of *Salmonella* genus

Genomic Island	<i>S. bongori</i> 12419	<i>S. arizonae</i> RKS2983	<i>S. typhimurium</i> LT2	<i>S. typhi</i> Ty2
SPI-1	+	+	+	+
SPI-2	-	+	+	+
SPI-3	+	-	+	+
SPI-4	+	+	+	+
SPI-5	+	+	+	+
SPI-6	-	-	+	+
SPI-7	-	-	-	+
SPI-8	-	-	-	+
SPI-9	+	+	+	+
SPI-10	-	-	-	+
SPI-11	+	+	+	+
SPI-12	-	-	+	+
SPI-13	+	+	+	-
SPI-14	-	+	+	-
SPI-15	-	-	-	+
SPI-16	-	-	+	+
SPI-17	-	-	-	+
SPI-18	-	-	-	+
SPI-19	-	-	-	-
SPI-20	+	+	-	-
SPI-21	+	+	-	-
SPI-22	-	-	-	-

+ means SPI is present in the serotype.

- means SPI is absent in the serotype.

The core gene data of *S. arizonae* RKS2983, *S. bongori* NCTC 12419 and *S. typhimurium* LT2 (representing *Salmonella* subgroup I) were presented in Figure 3. There are 2823 genes common to all three genomes and 926 genes specific in RKS2983. SPI-2 is in the set of 516 genes common to RKS2983 and LT2 and is absent in *S. bongori* NCTC 12419. As many as 1017 genes are in LT2 but not in the other two genomes; we postulate that some of these genes may be associated with virulence to warm-blooded hosts.

We compared these genomes for presence or absence of *Salmonella* pathogenicity islands (SPIs) and found that *S. arizonae* RKS2983 shared some of the SPIs with *S. bongori* NCTC 12419 and others with *S. typhimurium* LT2 or *S. typhi* Ty2 (Table 5), providing opportunities of evolutionary studies about acquisition of SPIs during transition of *Salmonella* from cold- to warm-blooded animal pathogens.

Conclusions

S. arizonae is phylogenetically positioned between *S. bongori* and *Salmonella* subgroup I and shares some pathogenicity-associated genes with *S. bongori* and some

others with *Salmonella* subgroup I lineages. Therefore *S. arizonae* genome analyses may provide important clues to key genomic events that might have facilitated the evolution of warm-blooded animal pathogens from cold-blooded parasites.

Abbreviations

SARC: *Salmonella* reference collection C; CTAB: Cetyl trimethyl ammonium bromide; MLEE: Multilocus enzyme electrophoresis.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

CXW carried out the genome sequence analysis and drafted the manuscript. SLZ and BL participated in genome sequence analysis. XYW participated in PCR amplification and sequencing of the PCR products by ABI3130 sequencer. RJ, JZ and GRL participated in the study design and provided reagents for the project. YGL and JZ provided the SOLiD and ABI Sequencing platform. SLL conceived the study and finalized the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by a Heilongjiang Innovation Endowment Award for graduate studies to CXW (YJSCX2012-235HLJ) and to XYW (YJSCX2012-198HLJ); National Natural Science Foundation of China (NSFC30970078 and NSFC81201248) and a grant of Natural Science Foundation of Heilongjiang Province of China to GRL; and grants of the National Natural Science Foundation of China (NSFC30970119, 81030029, 81271786, NSFC-NIH 81161120416) to SLL.

Author details

¹Genomics Research Center, Harbin Medical University, Harbin, China.

²Institute for Translational Medicine, Zhejiang University School of Medicine, Hangzhou, China. ³Department of Infectious Diseases, The First Affiliated Hospital, Harbin Medical University, Harbin, China. ⁴Department of Biochemistry and Molecular Biology, University of Calgary, Calgary, Canada.

⁵Department of Hematology, The First Affiliated Hospital, Harbin Medical University, Harbin, China. ⁶Department of Microbiology, Immunology and Infectious Diseases, University of Calgary, Calgary, Canada.

Received: 29 September 2014 Accepted: 21 April 2015

Published online: 03 June 2015

References

- Brenner FW, Villar RG, Angulo FJ, Tauxe R, Swaminathan B. *Salmonella* nomenclature. *J Clin Microbiol*. 2000;38(7):2465–7.
- Boyd EF, Wang FS, Whittam TS, Selander RK. Molecular genetic relationships of the salmonellae. *Appl Environ Microbiol*. 1996;62(3):804–8.
- Tang L, Wang CX, Zhu SL, Li Y, Deng X, Johnston RN, et al. Genetic boundaries to delineate the typhoid agent and other *Salmonella* serotypes into distinct natural lineages. *Genomics*. 2013;102(4):331–7.
- Tang L, Li Y, Deng X, Johnston RN, Liu GR, Liu SL. Defining natural species of bacteria: clear-cut genomic boundaries revealed by a turning point in nucleotide sequence divergence. *BMC Genomics*. 2013;14:489.
- Tang L, Liu SL. The 3Cs provide a novel concept of bacterial species: messages from the genome as illustrated by *Salmonella*. *Antonie Van Leeuwenhoek*. 2012;101(1):67–72.
- Baumler AJ, Tsois RM, Ficht TA, Adams LG. Evolution of host adaptation in *Salmonella enterica*. *Infect Immun*. 1998;66(10):4579–87.
- Doolittle RF, Feng DF, Tsang S, Cho G, Little E. Determining divergence times of the major kingdoms of living organisms with a protein clock. *Science (New York, NY)*. 1996;271(5248):470–7.
- Lawrence JG, Ochman H. Molecular archaeology of the *Escherichia coli* genome. *Proc Natl Acad Sci U S A*. 1998;95(16):9413–7.
- Porwollik S, McClelland M. Lateral gene transfer in *Salmonella*. *Microbes and infection / Institut Pasteur*. 2003;5(11):977–89.

10. Jones BD, Ghori N, Falkow S. *Salmonella typhimurium* initiates murine infection by penetrating and destroying the specialized epithelial M cells of the Peyer's patches. *J Exp Med*. 1994;180(1):15–23.
11. McCormick BA, Miller SI, Carnes D, Madara JL. Transepithelial signaling to neutrophils by salmonellae: a novel virulence mechanism for gastroenteritis. *Infect Immun*. 1995;63(6):2302–9.
12. Schmidt H, Hensel M. Pathogenicity islands in bacterial pathogenesis. *Clin Microbiol Rev*. 2004;17(1):14–56.
13. Ochman H, Groisman EA. Distribution of pathogenicity islands in *Salmonella* spp. *Infect Immun*. 1996;64(12):5410–2.
14. Hensel M, Shea JE, Baumlér AJ, Gleeson C, Blattner F, Holden DW. Analysis of the boundaries of *Salmonella* pathogenicity island 2 and the corresponding chromosomal region of *Escherichia coli* K-12. *J Bacteriol*. 1997;179(4):1105–11.
15. Hensel M, Shea JE, Waterman SR, Mundy R, Nikolaus T, Banks G, et al. Genes encoding putative effector proteins of the type III secretion system of *Salmonella* pathogenicity island 2 are required for bacterial virulence and proliferation in macrophages. *Mol Microbiol*. 1998;30(1):163–74.
16. Cirillo DM, Valdivia RH, Monack DM, Falkow S. Macrophage-dependent induction of the *Salmonella* pathogenicity island 2 type III secretion system and its role in intracellular survival. *Mol Microbiol*. 1998;30(1):175–88.
17. Fookes M, Schroeder GN, Langridge GC, Blondel CJ, Mammina C, Connor TR, et al. *Salmonella bongori* provides insights into the evolution of the Salmonellae. *PLoS Pathogens*. 2011;7(8):e1002191.
18. Ochman H, Soncini FC, Solomon F, Groisman EA. Identification of a pathogenicity island required for *Salmonella* survival in host cells. *Proc Natl Acad Sci U S A*. 1996;93(15):7800–4.
19. Crosa JH, Brenner DJ, Ewing WH, Falkow S. Molecular relationships among the Salmonellae. *J Bacteriol*. 1973;115(1):307–15.
20. SGC web site. [www.ucalgary.ca/~kesander]
21. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol*. 2008;26(5):541–7.
22. Doyle DJ. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull*. 1987;19:11–5.
23. Illumina web site. [http://www.illumina.com/technology/sequencing_technology.ilmn]
24. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics*. 2008;9:75.
25. Delcher AL, Bratke KA, Powers EC, Salzberg SL. Identifying bacterial genes and endosymbiont DNA with glimmer. *Bioinformatics (Oxford, England)*. 2007;23(6):673–9.
26. McClelland M, Sanderson KE, Spieth J, Clifton SW, Latreille P, Courtney L, et al. Complete genome sequence of *Salmonella enterica* serovar Typhimurium LT2. *Nature*. 2001;413(6858):852–6.
27. Desai PT, Porwollik S, Long F, Cheng P, Wollam A, Bhonagiri-Palsikar V, Hallsworth-Pepin K, Clifton SW, Weinstock GM, McClelland M: Evolutionary Genomics of *Salmonella enterica* Subspecies. *mBio*. 2013;4(2):e00198-13.
28. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res*. 1997;25(5):955–64.
29. Lagesen K, Hallin P, Rodland EA, Staerfeldt HH, Rognes T, Ussery DW. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res*. 2007;35(9):3100–8.
30. Griffiths-Jones S, Bateman A, Marshall M, Khanna A, Eddy SR. Rfam: an RNA family database. *Nucleic Acids Res*. 2003;31(1):439–41.
31. Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol*. 2001;305(3):567–80.
32. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res*. 2002;30(14):3059–66.
33. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol*. 2007;24(8):1596–9.
34. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci U S A*. 1990;87(12):4576–9.
35. Garrity GM, Bell JA, Lilburn T. Phylum XIV. Proteobacteria phyl. In Bergey's manual of systematic bacteriology, volume 2, part B. 2nd ed. New York: Springer; 2005:1.
36. Garrity GM, Bell JA, Lilburn T. Class III. Gammaproteobacteria class. In Bergey's manual of systematic bacteriology, volume 2, part B. 2nd ed. New York: Springer; 2005:1.
37. Garrity GM, Holt JG. Taxonomic outline of the archaea and bacteria. In Bergey's manual of systematic bacteriology, volume 1. 2nd ed. New York: Springer; 2001. p. 155–66.
38. Rahn O. New principles for the classification of bacteria. *Zentralbl Bakteriol Parasitenkd Infektionskr Hyg*. 1937;96:273–86.
39. Commission J. Conservation of the family name *Enterobacteriaceae*, of the name of the type genus, and designation of the type species Opinion No. 15. *Int Bull Bacteriol Nomencl Taxon*. 1958;8:74.
40. Gouillet P. Esterase electrophoretic pattern relatedness between *Shigella* species and *Escherichia coli*. *J Gen Microbiol*. 1980;117:493–500.
41. Tindall BJ, Grimont PA, Garrity GM, Euzéby JP. Nomenclature and taxonomy of the genus *Salmonella*. *Int J Syst Evol Microbiol*. 2005;55(Pt 1):521–4.
42. Le Minor L, Popoff MY. Request for an opinion. Designation of *Salmonella enterica* sp. nov., nom. rev., as the type and only species of the genus *Salmonella*. *Int J Syst Bacteriol*. 1987;37:465–8.
43. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. The gene ontology consortium. *Nat Genet*. 2000;25(1):25–9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

